

ED Treatment Variables:

Antiemetic medications administered while the patient was in the ED:

Dose 1: Type_____	Dose/Route_____	Time Administered_____
Dose 2: Type_____	Dose/Route_____	Time Administered_____
Dose 3: Type_____	Dose/Route_____	Time Administered_____
Dose 4: Type_____	Dose/Route_____	Time Administered_____
Dose 5: Type_____	Dose/Route_____	Time Administered_____
Dose 6: Type_____	Dose/Route_____	Time Administered_____
Dose 7: Type_____	Dose/Route_____	Time Administered_____
Dose 8: Type_____	Dose/Route_____	Time Administered_____

IV Fluid Therapy administered while the patient was in the ED:

IV Fluid 1: Type_____	Volume_____mL	Time Administered_____
IV Fluid 2: Type_____	Volume_____mL	Time Administered_____
IV Fluid 3: Type_____	Volume_____mL	Time Administered_____
IV Fluid 4: Type_____	Volume_____mL	Time Administered_____
IV Fluid 5: Type_____	Volume_____mL	Time Administered_____
IV Fluid 6: Type_____	Volume_____mL	Time Administered_____

Antiemetic prescriptions provided upon discharge from the ED:

Medication 1: Type_____	Dose/Route_____
Medication 2: Type_____	Dose/Route_____
Medication 3: Type_____	Dose/Route_____
Medication 4: Type_____	Dose/Route_____

Other Relevant Notes: